

INDEX

Synonyms are printed in *italics*; new names in **bold-face type**

- Acanthopanax aculeatum*, 71
— *trifoliatus*, 71
- Acer Buergerianum formosanum*, 216
— *trifidum formosanum*, 216
- Acicalyptus*, 326
— *elliptica*, 330
— *eugenoides*, 330
— *Fullageri*, 331
— *longiflora*, 329
— *myrtoides*, 329
— *nitida*, 337
— *Seemannii*, 330
- Acronychia Esquirolii*, 235
- Actinidia eriantha*, 222
— *lanata*, 222
- Adenosma glutinosum*, 73
- Adenosma grandiflorum*, 73
- Adina Fauriei*, 247
— *rubella*, 247
- Aeschynanthus* spec., 246
- Aganosma cymosa*, 237, 361
— *Schlechterianum*, 238
- Aglaia tetrapetala*, 210
- Akebia Cavaleriei*, 256
- Alchornea Mairei*, 214
- Alstonia Esquirolii*, 235
— *Mairei*, 235, 276
— *paupera*, 235
— *yunnanensis*, 235
- Alyxia Bodinieri*, 235
— *Schlechteri*, 235, 361
- Ampelopsis aconitifolia*, 220
— *cantonensis*, 71
— *Chaffanjoni*, 220
— *Delavayana Gentiliana*, 219
- Amsinckia intermedia*, 22
- Andrachne attenuata*, 212, 257
— *Cavaleriei*, 232
— *Esquirolii*, 212, 257, 361
— *Millietii*, 232
— *persicariifolia*, 257, 361
- Andrews, Henry C. Roses — Bibliographical Notes, 258
- Andropogon intermedius Haenkei*, 57
- Andropogon Vachellii*, 57
- Aneilema malabaricum*, 66
— *nudicaule*, 66
— *nudiflorum*, 65
- Antiotrema Dunnianum*, 25
- Antiphytum mexicanum*, 16
- Antirrhaea Esquirolii*, 235
— *Martini*, 238
- Aplos Martini*, 208, 276
- Aralia trifoliata*, 71
- Ardisia crispa*, 232
— *Dielsii*, 232
— *Henryi Dielsii*, 232
— *hortorum*, 232
- Arduina Carandas*, 235
- Argyreia Seguini*, 242
- Aristolochia Blinii*, 240
— *Mairei*, 240
— *viridiflora*, 240
— — *occlusa*, 240
- Arnebia Hancockiana*, 21
- Arnold Arboretum during the fiscal year ended June 30, 1937, The, 349
- Arnold Arboretum, Staff of the, 360
- Arundina chinensis*, 66
— *Meyenii*, 66
- Arundinella setosa*, 57
- Aspidopterys hypoglaucia*, 216
- Aster Cavaleriei*, 252
— *Fordii*, 74
— *lofouensis*, 252
— *lofouensis*, 277
— *panduratus*, 74
— *Walpersianus*, 74
- Atkins Institution of the Arnold Arboretum, 355
- BAILEY, I. W. and MARY R. VESTAL, The orientation of cellulose in the secondary wall of tracheary cells, 185, figs. pl. 206-208
- and MARY R. VESTAL, The significance of certain wood-destroying fungi in the study of the enzymatic hydrolysis of cellulose, 196, figs. pl. 209, 210

- BAILEY, I. W. and THOMAS KERR, The structural variability of the secondary wall as revealed by "lignin" residues, 261, pl. 211-214
- Bauhinia chinensis*, 68
— variegata, 68
- Benzoin commune*, 275
— *glaucum*, 275
— *grandifolium*, 275
— *touyunense*, 275
— — *megaphyllum*, 275
- Berghausia barbulata*, 58
- Bibliographical Notes, 183, 258
- Bibliography of the published writings of the staff and students, 357
- Bidens Meyeniana*, 76
— *tenuifolia*, 76
- Bileveillea granulatifolia*, 253
- Biophytum Esquirolii, 209
— *Reinwardtii*, 209
- Bischofia javanica, 213
- Blumea *amplectens*, 75
— *chinensis*, 75
— *conyzoides*, 252, 277
— *granulatifolia*, 253
— *hieracifolia*, 75
— *obliqua*, 75
— *lanceolaria*, 253
- Bodinierella Cavaleriei*, 230
- Boea Cavaleriei*, 246
- Boenninghausenia albiflora, 210
— — *brevipes*, 210
— — *sessilicarpa*, 210
- Boraginaceae, XII, Studies in the, 1
- Brackenridgea Hookeri, 204, pl. 209, fig. 5, pl. 210, fig. 8, 10, 11
- Brandisia racemosa, 245
- Bridelia spec., 213
- Buddleia *acutifolia*, 234
— — *albiflora*, 276
— — *Mairei albiflora*, 276
— — *officinalis*, 234
— — *albiflora*, 276
— *tibetica truncatifolia*, 234
— — *truncatifolia*, 234
- Burretiodendron Esquirolii, 220
- Buxus megistophylla, 215
— Myrica, 215
- Caeoma Empetri*, 145
- Callerya tomentosa*, 70
- Callicarpa Bodinieri, 243
— — *Giraldii*, 243
— — *Lyi*, 243
— — *Esquirolii*, 243
— — *Lyi*, 243
— — *Mairei*, 243
— — *Martini*, 243
- Calyptranthes*, 326
— *eugenoides*, 330
- Makal*, 337
- mangiferifolia*, 337
- myrtoides*, 329
- Seemanni*, 330
- Zuzygium*, 337
- Camellia Costei, 223
— *drupifera*, 223
- Grijsii*, 223
- japonica*, 223
- Mairei*, 223
- oleosa*, 223
- Pitardii*, 223
— — *lucidissima*, 223
- Camptotheca acuminata, 226, 361
- Campylotropis polyantha, 276
- Canthium Dunnianum*, 249
— *Labordei*, 249
- Capparis cantoniensis, 257
- Capraria crustacea*, 73
- Carissa Carandas, 235
- Carya aquatica* × *Pecan*, 133
— *cordiformis* × *ovalis*, 135
- Demareei*, 135
- texana*, 133
- Caryopteris *Mairei*, 244, 361
— *paniculata*, 243
- Castanopsis spec., 224
- Cayratia oligocarpa, 220
- Celastrus euonymoidea*, 221
— *Feddei*, 220, 221
- Celtis Biondii heterophylla, 255
— *Bungeana heterophylla*, 255
- Leveillei*, 255
- polycarpa*, 213
- Centrochilus gracilis*, 67
- Cephalanthus Esquirolii*, 226, 361
- Ceropegia Balfouriana, 240
- Chinese species described in Meyen's "Observationes Botanicae" (Beiträge

- zur Botanik), The, 54
Choeradoplectron spiranthes, 67
Chrysanthemum indicum, 253
 — — *coreanum*, 253
Chukrasia tabularis, 210
Chrysomyxa Empetri — A spruce-infecting rust, 141
Chrysomyxa Empetri, 145, fig., pl. 202
Cissampelos hypoglauca, 68
Cissus cantoniensis, 71
 — *diversifolia*, 71
Cladium chinense, 60
 — *jamaicense*, 60
 — *japonicum*, 60
Clausena Dunniana, 210
 — *excavata*, 210
 — *Willdenowii*, 210
Cleistocalyx Blume (including *Acicalyptus* A. Gray), a valid genus of the Myrtaceae, Reinstatement and revision of, 322, pl. 215
Cleistocalyx, 326, pl. 215
 — sect. **Acicalyptus**, 326
 — sect. **Eucleistocalyx**, 327
 — *arcuatinerius*, 333
 — *Baeurlenii*, 331
 — *barringtonioides*, 332
 — *Brongniartii*, 337
 — *conspersipunctatus*, 335
 — *ellipticus*, 330
 — *eugenoides*, 330
 — *Fullageri*, 331
 — *gustavioides*, 334
 — *leucocladus*, 336
 — *longiflorus*, 329
 — *myrtoides*, 329
 — *nervosus*, 342
 — *nicobaricus*, 341
 — *nigrans*, 336
 — *nitidus*, 333
 — *operculatus*, 337
 — *paradoxus*, 331
 — *paucipunctatus*, 336
 — *perspicuinervius*, 332
 — *retinervius*, 334
 — *Seemannii*, 330
 — — *punctatus*, 330
Clematis Finetiana, 256
 — *Meyeniana*, 67
 — *Pavoliniana*, 256
Clematoclethra spec., 222
Clethra Bodinieri, 228
 — *Cavaleriei*, 228
 — *kaipoensis*, 228
Cleyera, Studies in Theaceae, II, 118
Cleyera, 119
 — *albopunctata*, 128
 — *elegans*, 128
 — *Fortunei*, 123
 — *grandiflora*, 125
 — *integriifolia*, 128
 — *japonica*, 120
 — — *foliis variegatis*, 123
 — — *grandiflora*, 125
 — — *Hayatai*, 124
 — — *lipingensis*, 127
 — — *Morii*, 126
 — — *parvifolia*, 125
 — — *tricolor*, 123
 — — *tricolor*, 123
 — *Lushia*, 121
 — *Lushia Wallichiana*, 121
 — *Matsudai*, 128
 — *Mertensiana*, 121
 — *mexicana*, 128
 — *Nimanimae*, 128
 — *ochnacea*, 120
 — — *grandiflora*, 125
 — — *Kaempferiana*, 121
 — — *Lushia*, 121
 — *ochnoides*, 121
 — *serrulata*, 128
 — *siphilitica*, 129
 — *theoides*, 129
 — *Wallichiana*, 121
 Climate and vegetation in southern New England and adjacent New York, Recent changes of, 79
Clitoria ternatea, 68
Commelinia cespitosa, 64
 — *diffusa*, 64
 — *longicaulis*, 64
 — *nudicaulis*, 66
 — *nudiflora*, 64, 66
 — *ochreata*, 64
 — *pacifica*, 64
 — *trichocolea*, 66
Compsoneura capitellata, 204, pl. 209, fig. 3
 Control of the Gymnosporangium rusts

- by means of sulphur sprays, 149
Conyza amplexicaulis, 75
 — *Cappa*, 76
 — *obliqua*, 75
 — *pyrifolia*, 77
 — *syringaeifolia*, 77
 — *viscidula*, 252, 277
 — *volubilis*, 77
Cordia cordiformis, 10
 — *decipiens*, 11
 — *lomatoloba*, 12
 — *Mexiana*, 12
Cornus florida, 344
CREAGER, D. B., Phytophthora crown rot of dogwood, 344, pl. 216
Crotalaria albida, 68
 — *elliptica*, 68
 — *leiocarpos*, 68
 — *Mairei*, 206
 — *splendens*, 68
 — *Vachellii*, 68
 — *spec.*, 211
CROWELL, IVAN H. and J. D. MACLACHLAN, Control of the Gymnosporangium rusts by means of sulphur sprays, 149
Cryptantha corollata, 24
 — *decipiens corollata*, 24
 — *Grahamii*, 23
 — *Hooveri*, 23
Cryptolepis Buchananii, 239
Cudrania Bodinieri, 257
 — *tricuspidata*, 275
Cyclea deltoidea, 68
 — *hypoglaucia*, 67
Cymbidium ensifolium, 66
 — *Meyenii*, 66
 — *micans*, 66
Cynanchum Boudieri, 239
 — *caudatum*, 239
 — *Mooreanum*, 239
 — *tylophoroideum*, 239
 — *yunnanense*, 249
Cynoglossum Cavaleriei, 25
 — *Dunnianum*, 25
Cyperus compressus, 61
 — *Griffithii*, 61
 — *Meyenii*, 61
 — *radians*, 61
 — *radicans*, 61
Cyperus sinensis, 61
Dalbergia Cavaleriei, 208
Damnacanthus Esquirolii, 235
Daphne Bodinieri, 235
 — *Feddei*, 226
 — *Martini*, 226
Davidia involucrata, 204, pl. 209, fig. 7
Dendropanax morbiferus, 227
Desmodium cinerascens, 207
 — *Esquirolii*, 207
 — *formosum*, 69
 — *Franchetii*, 207
 — *heterocarpum*, 69
 — *nervosum*, 69
 — *polycarpum*, 69
 — *viride*, 70
Deutzia funebris, 245
Didymocarpus Cavaleriei, 246
Digitalis sinensis, 73
Digitaria dispar, 58
 — *heterantha*, 58
Diospyros Bodinieri, 215
 — *Esquirolii*, 233
Diploid and polyploid plants, Stomata size and distribution in, 164
Dischidia Esquirolii, 241
 — *yunnanensis*, 241
Dogwood, Phytophthora crown rot of, 344, pl. 216
Dolichos trilobus, 209
Duhaldea chinensis, 76
Dumasia hirsuta, 208
 — *villosa*, 208, 276
Ecdysanthera rosea, 235
Echium connatum, 21, 250, 276
Edwards's Botanical Register — Bibliographical Notes, 183
Ehretia acuminata, 243
 — *Argyi*, 243
Elaeagnus Argyi, 226
 — *coreanus*, 226
 — *crispa coreana*, 226
 — — *parvifolia*, 226
 — *umbellata*, 226
Elsholtzia Bodinieri lavandulaespica, 244
 — *heterophylla*, 244, 361
 — *lavandulaespica*, 244
Embelia Bodinieri, 232
 — *oblongifolia*, 232

- Embelia rubrinervis*, 219
Enkianthus Cavaleriei, 230
 — *cerasiflora*, 230
 — *chinensis*, 230
 — *Dunnii*, 230
 — *xanthoxanthus*, 230
Epidendrum ensifolium, 66
Eragrostis cylindrica, 59
 — *geniculata*, 59
 — *Millettii*, 59
 — *pilosissima*, 59
Erigeron hieracifolium, 75
 — *obliquum*, 75
Eriolaena Esquirolii, 220
Eriosema chinense, 69
Erithalis polygama, 131
 — *Timon*, 131
Eritrichium brevipes, 6
 — *microcarpum*, 8
 — *pedunculare*, 8
Erysibe Empetri, 145
Eugenia, 326
Eugenia arcuatina, 333
 — *Baeurlenii*, 332
 — *barringtonioides*, 332
 — *clausa*, 338
 — *cleistocalyx*, 333
 — *divaricato-cymosa*, 338
 — *gustavioides*, 334
 — *Holteana*, 338
 — *Holtzeana*, 338
 — *Holtzei*, 337
 — *microphylla*, 71
 — *nervosa*, 342
 — *nervosa*, 333
 — *nicobarica*, 341
 — *nigrans*, 336
 — *occlusa*, 341, 342
 — *operculata*, 337
 — *paradoxa*, 331
 — *paucipunctata*, 336
 — *perspicuinervia*, 332
 — *prora*, 330
 — *sinensis*, 71
 — *subrufa*, 336
 — *sympysipetala*, 342
Euploca racemosa, 19
Eurya § *Cleyera*, 120
 — subg. *Cleyera*, 120
 — *Esquirolii*, 290
 — *latifolia variegata*, 123
 — *Matsudai*, 128, 129
 — *ochnacea*, 121
 — — *lipingensis*, 127
 — — *Morii*, 126
Eurycorymbus austrosinensis, 217
 — *Cavaleriei*, 217
Evonymus centidens, 215
 — *Thunbergiana oblongifolia*, 215
Fagara Chaffanjoni, 210
 — *odorata*, 209
FAULL, J. H., *Chrysomyxa Empetri* —
 A spruce-infecting rust, 141, fig., pl.
 202, 203
Ferula marathrophylla, 72
Ficus corymbifera, 245
 — *Esquirolii*, 256
 — *foveolata*, 255
 — *hirtaeformis*, 222
 — *Martini*, 256
 — *Ouangliense*, 211
 — *rufipes*, 248, 255
 — *Vanioti*, 256
 — *Vanioti*, 211
Fimbristylis decora, 62
 — *podocarpa*, 62
 — *sericea*, 62
 — *subbispicata*, 61
Fraxinus Fauriei, 217
 — *mandshurica*, 195, pl. 208, fig. 14
Freziera elegans, 128
 — *integrifolia*, 128
 — *Nimanimae*, 128
 — *ochnacea*, 121
 — *theoides*, 129
 — sp., 128
Fusarium Caries, 56
Garnotia barbulata, 58
Gerardia glutinosa, 73
Gilibertia morbifera, 227
Glossogyne tenuifolia, 76
Glycine Soja, 208
 — *ussuriensis*, 208
Gongronema yunnanense, 242
Gordonia axillaris, 224
Grewia biloba, 221
 — *Esquirolii*, 221
 — *Feddei*, 220

- Grewia glabrescens*, 221
 — *Henryi*, 221
 — *parviflora glabrescens*, 221
 — *spec.*, 221
Gymnosporangium rusts by means of sulphur sprays, Control of the, 149
Habenaria acuifera, 66
 — *endothrix*, 67
 — *lacertifera*, 67
 — *linguella*, 66
 — *Meyenii*, 67
Hackelia brachytuba, 25
 — *Dielsii*, 25
Handeliodendron Bodinieri, 217
Hedera hypoglauca, 71
Hedyotis Esquirolii, 247
 — *hedyotidea*, 247
Hedysarum heterocarpon, 69
 — *lutescens*, 70
 — *polycarpon*, 69
 — *sericeum*, 70
Heliotropium americanum, 17
 — *angustifolium*, 18
 — *anomalum mediale*, 21
 — *calcicola*, 16
 — *convolvulaceum racemosum*, 19
 — *coriaceum*, 15
 — *dichroum*, 18
 — *Eggersii*, 17
 — *eremogenum*, 20
 — *fallax*, 14
 — — *Hintonii*, 15
 — *inundatum*, 17
 — — *cubense*, 17
 — *Jaffuelii*, 19
 — *petraeum*, 17
 — *procumbens*, 17
 — *pueblense*, 17
 — *rigidulum*, 17
 — *Sessei*, 13, 361
 — *uninerve*, 18
Henreyettana mirabilis, 25
Herbarium, On the technique of inserting published data in the, 173
Heterostemma Esquirolii, 242
Hoya carnosa, 241
 — *Esquirolii*, 241
 — *Lyi*, 241
Hoyopsis Dielsii, 241
Hypericum Bodinieri, 225
 — *Dielsii*, 224
 — *erectum*, 224
 — — *axillare*, 224
 — *Matsumurae*, 224
 — *napaulense*, 225
 — *otaruense*, 224
 — *Vanioti*, 224
Ilex macrocarpa, 215
 — *purpurea*, 215
Illigera Dunniana, 256
Incarvillea Argyi, 242
Index to Notes on the ligneous plants described by H. Léveillé from Eastern Asia, 278
Indigofera chinensis, 70
 — *Esquirolii*, 207
Inula Cappa, 76
 — *indica hypoleuca*, 277
 — *pseudo-cappa*, 76
Iodes ovalis, 276
Iryanthera macrophylla, 204, pl. 209, fig. 4
Ischaemum aristatum lodiculare, 59
 — — *Meyenianum*, 59
Ixora Henryi, 248
Jambosa nitida, 333
Jasminum Prainii, 234
 — *Seguini*, 233
JOHNSTON, IVAN M., Studies in the Boraginaceae, XII, 1
KERR, THOMAS, and *I. W. BAILEY*, The structural variability of the secondary wall as revealed by "lignin" residues, 261, pl. 211-214
Keteleeria Davidiana, 254
 — *Esquirolii*, 254
KOBUSKI, CLARENCE E. Studies in Theaceae. II, *Cleyera*, 118, pl. 201
Lappula Dielsii, 25
Larix occidentalis, 195, 271, pl. 207, fig. 5, 7, 8, pl. 214, fig. 14
Lasianthus Hartii, 249
 — *Labordei*, 249
Laurelia aromatica, 204, pl. 209, fig. 6, pl. 210, fig. 9
Lepiaglaia ? tetrapetala, 210
Lepidosperma chinense, 62
Lespedeza dichromocalyx, 276

- Lespedeza dichromoxylon*, 276
 — *fasciculiflora*, 207
 — *formosa*, 69
 — *Monnoyeri*, 208
 — *sericea*, 69
 — *viatorum*, 69
Lettsomia Seguini, 242
Leucothoë Griffithiana, 230
 — *spec.*, 230
 Léveillé, Notes on the ligneous plants described by H. . . . from Eastern Asia, 26, 206, 273
 Lignin residues, The structural variability of the secondary wall as revealed by, 261, pl. 211-214
Lindera Bodinieri, 275
 — *communis*, 275
 — *glauca*, 275
 — *megaphylla*, 275
 — — *touyunensis*, 275
 — *yunnanensis*, 275
 — *spec.*, 213
Lindernia crustacea, 73
Lithocarpus edulis, 195, pl. 208, fig. 12
Lithodora Hancockiana, 21
Lithospermum albiflorum, 21
 — *Hancockianum*, 21
 — *Komarovianum*, 22
 — *Mairei*, 21
 — *officinale*, 21
Litsaea Esquirolii, 275
Litsea Cavaleriei, 275
 — *Kobuskiana*, 290
 — *touyunensis*, 275
Lonicera Pampaninii, 250, 361
Lophopetalum spec., 271, pl. 213, fig. 9
Lycopodium amentigerum, 57
 — *cernuum*, 57
Lyonia ovalifolia, 230
 — — *lanceolata*, 231
 — *villosa pubescens*, 231
Lysimachia capillipes Cavaleriei, 232
 — *Mairei*, 222
 — *Millietii*, 232
Lysionotus Cavaleriei, 246
 — *pauciflorus linearis*, 246
Macaranga Esquirolii, 214
 MACLACHLAN, J. D. and IVAN H. CROWELL, Control of the Gymnosporangium rusts by means of sulphur sprays, 149, pl. 204
Maesa scandens, 237
Mallotus Esquirolii, 213
Mariscus chinensis, 60
Marquartia tomentosa, 70
Marsdenia Cavaleriei, 242
 — *oreophila*, 242
 — *stenantha*, 241
 — *tenacissima*, 242
 — *yunnanensis*, 242
Meibomia viridis, 70
Melampsoropsis Empetri, 145
Melica latifolia, 60
Meliosma cuneifolia, 218
 — *Oldhamii*, 217
Melodinus Bodinieri, 237
 — *Cavaleriei*, 236, 237
 — *Chaffanjoni*, 237
 — *Dunnii*, 237
 — *Hemsleyanus*, 234, 361
 — *khasianus*, 234
 — *Seguini*, 235
Melodium Dunnii, 237
Meoschium lodiculare, 59
 — *Meyenianum*, 59
 MERRILL, E. D., The Chinese species described in Meyen's "Observationes Botanicae" (Beiträge zur Botanik), 54
 — New names in Timonius, 130
 — On the technique of inserting published data in the herbarium, 173
 — and L. M. PERRY, Reinstatement and revision of Cleistocalyx Blume (including Acicalyptus A. Gray), a valid genus of the Myrtaceae, 322, pl. 215
Mespilus Esquirolii, 222
Metaplexis Cavaleriei, 242
 Meyen's "Observationes Botanicae" (Beiträge zur Botanik), The Chinese species described in, 54
Microglossa albescens, 252
 — *pyrifolia*, 77
 — *volubilis*, 77
Microtoena Esquirolii, 244
 — *insuavis*, 244
 — *mollis*, 244
Microtropis fokienensis, 216
Milletia Esquirolii, 207

- Milletia nitida*, 70
Miquelia barbulata, 58
Mitrasacme chinensis, 72
 — *nudicaulis*, 72
Mitreola pedicellata, 10
Moquinia eriosematooides, 76
Morinda Esquirolii, 214
Morus integrifolia, 275
Mucuna Birdwoodiana, 208
 — *Bodinieri*, 208
Mussaenda Esquirolii, 248
Myodocarpus simplicifolius, 195, pl. 208,
 fig. 10, 11
Myosotis mexicana, 14
 — *microcarpa*, 8
 — *peduncularis*, 8
Myrsine Chaffanjoni, 216
Myrtus Loureiri, 342
Nelitris Timon, 131
Neolitsea spec., 290
 New names in *Timonius*, 130
 North American trees and shrubs, Notes
 on, 133, figs.
 Notes on the ligneous plants described by
 H. Léveillé from Eastern Asia, 26, 206,
 273
 Notes on North American trees and
 shrubs, 133, 2 figs.
Oldenlandia corymbosa, 73
 — *hedyotidea*, 247
 — *macrostemon*, 247
Omphalodes Bodinieri, 10
 — *Cavaleriei*, 6
 — *cordata*, 5
 — *Esquirolii*, 6
 — *Mairei*, 4
 — *moupinensis*, 5
 — *sericea*, 25
 — *Vaniotii*, 6
Onosmodium dodrantale, 22
Ophiorrhiza Esquirolii, 234
 Orientation of cellulose in the secondary
 wall of tracheary cells, The, 185, pl.
 206-208, 3 figs.
Osteophloeum platyspermum, 205, pl.
 210, fig. 13
Paederia Bodinieri, 249
 — *Cavaleriei*, 249
 — *Dunniana*, 249
Paederia Esquirolii, 249
 — *Mairei*, 249
 — *Rehderiana*, 249
 — *scandens*, 249
 — *tomentosa purpureo-coerulea*, 249
 — *Wallichii*, 249
 — *yunnanensis*, 249
 PALMER, ERNEST J., Notes on North
 American trees and shrubs, 133, 2 figs.
Panax aculeatum, 71
Panicum acariferum, 60
 — *commutatum*, 58
 — *heteranthum*, 58
Paracaryum brachytubum, 25
Paradombeya sinensis, 222
Parameria Esquirolii, 238
Paspalum heteranthum, 58
Passiflora cupiformis, 225
 — Seguini, 225
Pentace Esquirolii, 220
Pentasacme Esquirolii, 242
Periploca astacus, 237
Peristylus chloranthus, 67
 PERRY, L. M. and E. D. MERRILL, Rein-
 statement and revision of *Cleistocalyx*
 Blume (including *Acicalyptus* A.
 Gray), a valid genus of the Myrta-
 ceae, 322, pl. 215
Phyllagathis Cavaleriei, 227, 361
Phyllanthus Bodinieri, 212
 — *spec.*, 212
Phytophthora cactorum, 345
Phytophthora crown rot of dogwood,
 344, pl. 216
Pieris Cavaleriei, 230
 — *Duclouxii*, 231
 — *Henryi*, 231
 — *kouyangensis*, 231
 — *longicornu*, 231
 — *Mairei*, 231
 — — *parvifolia*, 231
 — *repens*, 231
 — *Ulrichii*, 230
Pinus longifolia, 195, 272, pl. 208, fig. 9,
 pl. 214, fig. 15, 16
 — *rigida serotina*, 205, pl. 210, fig. 12
 — *Strobus*, 195, pl. 206, fig. 4
Piptanthus spec., 211
Pirus brunnea, 275

- Poa cylindrica*, 59
Podocarpus Mairei, 254
Pogonatherum paniceum, 60
— *refractum*, 60
— *saccharoideum*, 60
Pogostemon lavandulaespica, 244, 361
Polygala arillata, 211
— *Dunniana*, 211
— *japonica*, 211
— *Taquetii*, 211
Polyphragmon sericeum, 131
Populus adenopoda, 273
— *Bonatii*, 273
— *rotundifolia macranthela*, 273
Poraqueiba sericea, 204, 271, pl. 209, fig. 1, pl. 212, fig. 3-7
Premna Bodinieri, 243
— *Mairei*, 218
— *puberula*, 243
Prismatomeris Labordei, 249
Psychotria Henryi, 248
— *Prainii*, 248
Pteris ensiformis, 57
Pterostigma grandiflorum, 73
— *rubiginosum*, 73
Pterostyrax hispidus, 233
— *Leveillei*, 233
Pueraria caerulea, 276
— *coerulea*, 276
— *Thomsoni*, 209
— *Thunbergiana*, 208, 276
— *triloba*, 208
Pycnospora lutescens, 70
Quamoclit pennata, 242
Quercus annulata, 138
— *bicolor* × *Q. lyrata*, 140
— *breviloba*, 138
— *breviloba* × *Q. stellata*, 139
— *coreana*, 274
— *glandulifera*, 274
— *humidicola*, 140
— *Mahloni*, 139, 361
— *Nuttallii*, 136, fig. 1
— — *cachensis*, 136, fig. 2
— *prinoides* × *Q. stellata*, 139
— *serrata*, 274
— *stelloides*, 139
Ramalina digitata, 56
— *geniculata*, 56

RAUP, HUGH M., Recent changes of climate and vegetation in southern New England and adjacent New York, 79
Recent changes of climate and vegetation in southern New England and adjacent New York, 79
REHDER, ALFRED, Index to Notes on the ligneous plants described by H. Léveillé from Eastern Asia, 278
— Notes on the ligneous plants described by H. Léveillé from Eastern Asia, 26, 206, 273
Reinstatement and revision of *Cleistocalyx* Blume (including *Acicalyptus* A. Gray), a valid genus of the Myrtaceae, 322, pl. 215
Rhabdothamnopsis chinensis, 246
— *sinensis*, 246
Rhamnella Martini, 219
— *rubrinervis*, 219
Rhamnus Bodinieri, 219
— *Cavaleriei*, 219
— *Esquirolii*, 219
— *Leveilleanus*, 219
— *Martini*, 219
Rhizophora Mangle, 271, pl. 211, fig. 1
Rhynchosia Argyi, 208
Rhynchospora chinensis, 62
Rhododendron Bachii, 229
— *caeruleum*, 229, 361
— *chrysocalyx*, 230
— *denudatum*, 229
— *Duclouxii*, 229
— *Esquirolii*, 229
— *irroratum*, 229, 361
— *kouytchense*, 230
— *Maximowiczianum*, 229, 361
— *rex*, 229
— *spinuliferum*, 229
— *Vaniotii*, 229
Rosa acicularis, 257
— *Argyi*, 257
— *cymosa*, 257
— *korsakoviensis*, 257
— *lucidissima*, 257
— *Marrettii*, 257
— *microcarpa*, 257
Rubus adenochlamys, 49
— *alceaefolius*, 32

- Rubus alnifoliolatus*, 41
 — *ampelophyllus*, 39
 — *amphidasys*, 35
 — *ampliflorus*, 30
 — *andropogon*, 33
 — *arbor*, 28
 — *Argyi*, 44
 — *asper*, 43
 — — *grandifoliolatus*, 43
 — — *myriadenus*, 43
 — *Blinii*, 37
 — *Bodinieri*, 37
 — *Bonatii*, 48
 — *Boudieri*, 49
 — *Buergeri*, 37
 — *calycacanthus*, 31
 — — *Buergerifolia*, 31
 — *Cavaleriei*, 34
 — *Chaffanjoni*, 35
 — *chroosepalus*, 28
 — *clinocephalus*, 33
 — *corchorifolius*, 37
 — *coreanus*, 46
 — — *Nakaianus*, 46
 — *crataegifolius*, 38
 — — *itoensis*, 39
 — — *Makinoensis*, 39
 — — *minor*, 39
 — — *subcrataegifolius*, 40
 — *croeacanthus*, 42
 — *Darrisii*, 31
 — *defensus*, 27
 — *Delavayi*, 41
 — *diamantinus*, 52
 — *Duclouxii*, 41
 — *eous*, 50
 — *erectifolius*, 39
 — *Esquirolii*, 34
 — *euleucus*, 50
 — *fasciculatus*, 50
 — *Fauriei*, 38
 — *Feddei*, 35
 — *fimbriiferus*, 32
 — *foliolosus*, 48
 — *fragariooides adenophora*, 26
 — *Franchetianus*, 26
 — *Gentilianus*, 29
 — *Grossularia*, 40
 — *hastifolius*, 36
- Rubus hirsutus*, 44
 — — *Argyi*, 44
 — *Hoatiensis*, 47
 — *holadenus*, 31
 — *ichangensis*, 29
 — *idaeus aculeatissimus*, 53
 — — *diamantinus*, 52
 — — *Matsumuranus*, 52
 — — *microphyllus*, 52
 — — *sachalinensis*, 53
 — — *strigosus*, 52
 — — *Yabei*, 51
 — *ikenoensis*, 27
 — *illudens*, 48
 — *incisus*, 40
 — — *geifolius*, 40
 — — *Koehneanus*, 41
 — — *subcrataegifolius*, 40
 — *innominatus Kuntzeanus*, 49
 — *irenaeus*, 36
 — *itoensis*, 39
 — *Jamini*, 36
 — *japonicus*, 26
 — *kanayamensis*, 51
 — *kerriifolius*, 38
 — *Kinashii*, 50
 — — *coreensis*, 49
 — *Koehneanus*, 41
 — *Kuntzeanus*, 49
 — — *xanthacantha*, 49
 — *Labbei*, 31
 — *Lambertianus minimiflorus*, 30
 — — *paykouangensis*, 30
 — *lasiocarpus micranthus*, 45
 — *longistylus* 49
 — *Lyi*, 35
 — *Mairei*, 28
 — *Mairei*, 48
 — *Makinoensis*, 39
 — *malifolius*, 28
 — *mallodes*, 33
 — *marmoratus*, 42
 — *Matsumuranus*, 52
 — *melanolasius discolor*, 53
 — *mesogaeus*, 50
 — *micranthus*, 45
 — *minimiflorus*, 30
 — *minusculus*, 41
 — *Mongouilloni*, 32

Rubus Mouyousensis, 28
 — *multibracteatus*, 33
 — — *Demangei*, 32
 — *myriadenus*, 43
 — — *grandifoliolatus*, 43
 — *Nakaianus*, 47
 — *nipponicus*, 51
 — *ouensanensis*, 40
 — *Papyrus*, 29
 — *parvifolius*, 47
 — *paykouangensis*, 30
 — *petaloideus*, 28
 — *pinfaensis*, 50
 — *pseudo-japonicus*, 26
 — — *diversifolius*, 26
 — *pseudosaxatilis*, 46
 — — *Kouytchensis*, 46
 — *Pyi*, 45
 — *quelpaertensis*, 46
 — *refractus*, 27
 — *Rocheri*, 27
 — *rosaefolius minor*, 42
 — — *sumatranus*, 43
 — *sachalinensis*, 53
 — *schizostylus*, 48
 — *setchuenensis*, 34
 — *sino-Sudrei*, 36
 — *stephanandria*, 44
 — *strigosus*, 52, 53
 — — *kanayamensis*, 51
 — *subcrataegifolius*, 41
 — *suberectifolius*, 39
 — *succedaneus*, 42
 — *sumatrarus*, 43
 — *talaikiaensis*, 44
 — *Taquetii*, 47
 — *tephrodes ampliflorus*, 30
 — *Thunbergii*, 44
 — — *Argyi*, 44
 — — *R. Argyi*, 44
 — — *talaikiensis*, 44
 — — *R. talaikiensis*, 44
 — *tongchouanensis*, 45
 — *triflorus diversifolius*, 26
 — *triphyllus*, 47
 — — *Taquetii*, 47
 — *Vanioti*, 38
 — *Wrightii ampelophyllus*, 39
 — — *makinoensis*, 39

Rubus Wrightii ouensanensis, 40
 — *xanthacantha*, 49
 — *Yabei*, 51
Saccharum paniceum, 60
Sakakia, 120
 — *Hayatai*, 124
 — *longicarpa*, 127
 — *Matsudai*, 129
 — *Morii*, 126
 — *ochnacea*, 121
Salix amygdalina nipponica, 273
 — *andropogon*, 225, 361
 — *angiolepis*, 255
 — *anisandra*, 254
 — *Argyi*, 254
 — *Blinii*, 274
 — *Camusii*, 254
 — *caprea*, 273
 — *Cavaleriei*, 254
 — *dodecandra*, 254
 — *Duclouxii*, 255
 — — *kouytchensis*, 255
 — *erioclada*, 255
 — *Feddei*, 273
 — *funebris*, 255
 — *Gilgiana*, 274
 — *gymnolepis*, 274
 — *hallaisanensis*, 273
 — — *nervosa*, 274
 — *Kinashii*, 273
 — *koreensis*, 273
 — *kouytchensis*, 255
 — *luctuosa*, 255
 — *Mairei*, 255
 — *Makinoana*, 274
 — *pachyclada*, 255
 — *polygonandra*, 273
 — *polyandra*, 254
 — *pseudo-Gilgiana*, 273
 — *pseudo-jessoensis*, 273
 — *pseudo-lasiogynne*, 273
 — *purpurea gymnolepis*, 274
 — *Pyi*, 254
 — *Taquetii*, 274
 — *triandra discolor*, 273
 — *variegata*, 255
 — *Wallichiana*, 255
 — *Wilsonii*, 254
 — *yunnanensis*, 254

- Sarcopyramis Bodinieri*, 227
 — *napalensis*, 227
 — *nepalensis Bodinieri*, 227
- SAX, HALLY JOLIVETTE and KARL SAX,
 Stomata size and distribution in diploid and polyploid plants, 164, pl. 205
- SAX, KARL and HALLY JOLIVETTE SAX,
 Stomata size and distribution in diploid and polyploid plants, 164, pl. 205
- Schistocodon Meyenii*, 72
- Scleria pratensis*, 63
 — *pterota*, 63
- Scoparia gypsophiloides*, 73
- Secondary wall as revealed by "lignin" residues, The structural variability of the, 261, pl. 211-214
- Secondary wall of tracheary cells, The orientation of cellulose in the, 185, pl. 206-208, figs.
- Senecio araneosus*, 253
 — *Gentilianus*, 254
 — *Hoi*, 277
 — *saxatilis*, 253
 — *spelaeicola*, 277
 — *Walkeri*, 253, 361
 — *Walkeri*, 277
 — *yalungensis*, 277
- Sequoia sempervirens*, 194, pl. 206, 208, figs. 1-3, 13
- Sideroxylon Bodinieri*, 217
- Significance of certain wood-destroying fungi in the study of the enzymatic hydrolysis of cellulose, The, 196, pl. 209, 210, figs.
- Sindechites Esquirolii*, 238
 — *Henryi*, 238
- Siparuna bifida*, 271, pl. 213, fig. 8, 10-13
- Solanum Capsicastrum*, 245
 — *cornutum*, 245, 361
 — *Dunnianum*, 245
 — *Heudesii*, 245, 361
 — *verbascifolium*, 245
- Sophora* spec., 207
- Spruce-infesting rust — *Chrysomyxa Empetri*, A, 141, pl. 202, 203
- Staff of the Arnold Arboretum, 360
- Stauntonia obovata*, 256
- Stemonurus secundiflorus*, 204, pl. 209, fig. 2
- Stephanotis yunnanensis*, 242
- Sterculia Bodinieri*, 212
- Stomata size and distribution in diploid and polyploid plants, 164
- Strobilanthes lofouensis*, 247
- Structural variability of the secondary wall as revealed by "lignin" residues, The, 261, pl. 211-214
- Strychnos Esquirolii*, 218
- Studies in the Boraginaceae, XII, 1
- Studies in Theaceae. II, *Cleyera*, 118
- Styrax Argyi*, 233
 — *Bodinieri*, 233
 — *Cavaleriei*, 233
 — *grandiflorus*, 233
 — *japonicus*, 233
 — *touchanensis*, 233
- Syllsygium buxifolium*, 71
- Symphitum fruticosum*, 16
- Symplocos coronigera*, 233
 — *Courtoisii*, 215
 — *Ernesti*, 233
- Syzygium* sect. *Symphytum*, 326
 — *angkolanum*, 337
 — *buxifolium*, 71
 — *costatum*, 341
 — *firmum*, 340
 — *fruticosum*, 342
 — *javanicum*, 342
 — *nervosum*, 337
 — *nodosum*, 337
 — *occlusum*, 342
 — *operculatum*, 338
- Tarenna incerta*, 248
- Technique of inserting published data in the herbarium, On the, 173
- Tephrosia vestita*, 71
- Ternstroemia albopunctata*, 128
 — *japonica*, 120
 — *Lushia*, 121
 — sp., 128, 129
- Tetramerista glabra*, 271, pl. 211, fig. 1
- Tetrastigma obtectum*, 219
- Teucrium palmatum*, 244, 361
- Textoria morbifera*, 228
- Thea Camellia lucidissima*, 223
 — *Cavaleriana*, 223
 — *Costei*, 223
 — *Grijsii*, 223

- Thea Mairei*, 223
 — *oleosa*, 223
 — *Pitardii*, 223
 — — *lucidissima*, 223
 — *podogyna*, 223
 — *sinensis androxantha*, 223
- Theaceae, Studies in, II, *Cleyera*, 118
- Thecopsozma Empetri*, 145
- Thysanolaena acarifera*, 60
 — *Agrostis*, 60
 — *maxima*, 60
 — *procera*, 60
- Timonius*, New names in, 130
- Timonius*, 131
 — *ferrugineus*, 132
 — *hirsutus*, 132
 — *involucratus*, 132
 — *octonervius*, 132
 — *papuanus*, 132
 — *Ridleyi*, 132
 — *Rumphii*, 131
 — *sericeus*, 131
 — *Timon*, 131
- Toxocarpus Wightianus*, 72
- Trachelospermum axillare*, 237, 361
 — *Bodinieri*, 237, 361
 — *Cavaleriei*, 239
 — *Dunnii*, 237, 361
 — *Esquirolii*, 234, 361
 — *gracilipes*, 236
 — — *Cavaleriei*, 236
 — *Navillei*, 238
 — *rubrinerve*, 236, 237
- Tradescantia malabarica*, 66
- Tragia involucrata*, 214
 — *Mairei*, 214
- Trigonotis*, 1
 — *Bodinieri*, 10
 — *brevipes*, 6
 — *Cavaleriei*, 6
 — *compressa*, 4
 — *contortipes*, 10
 — *delicatula*, 10
 — *Faberi*, 6
 — *gracilipes*, 9
 — *heliotropifolia*, 9
 — *macrophylla*, 4
 — — *trichocarpa*, 4
 — — *verrucosa*, 4
- Trigonotis Mairei*, 4
 — *microcarpa*, 7
 — *mollis*, 7
 — *moupinensis*, 5
 — *muriculata*, 4
 — *omeiensis*, 6
 — *peduncularis*, 8
 — — *macrophylla*, 4, 361
 — — *microcarpa*, 8
 — — *vestita*, 8, 361
 — *Rockii*, 10
 — *rotundata*, 7
 — *sericea*, 25
 — *vestita*, 8
- Triosteum himalayanum*, 21, 250, 276
- Tristylium*, 120
 — *ochnaceum*, 121
 — — *Morii*, 126
- Trochodendron aralioides*, 195, pl. 207, fig. 6
- Trypterygium hypoglauicum*, 216
- TUCKER, ETHELYN M., Bibliographical Notes, 183, 258
- Tylophora Argyi*, 239, 240
 — *Cavaleriei*, 239
 — *Dielsii*, 241
 — *floribunda*, 240
 — *Hoyopsis*, 241
 — *Leveilleana*, 240
 — spec. ?, 241
- Uredo Empetri*, 145
- Urena diversifolia*, 71
 — *lobata*, 71
- Utricularia racemosa*, 73
- Vaccinium Donianum*, 231
 — *Duclouxii*, 231
 — *fragile myrtifolium*, 231
 — *Mairei*, 231
 — *mandarinorum*, 231, 361
 — *mandarinorum austrosinense*, 231, 361
 — *repens*, 231
- Vandellia crustacea*, 73
 — *limosa*, 73
- Vanieria Bodinieri*, 257
 — *tricuspidata*, 275
- Vernonia arbor*, 277
 — *arbor*, 251
 — *arborea*, 250, 276
 — *congesta*, 76

- Vernonia eriosematoides*, 76
— *Esquirolii*, 251, 253, 277, 361
— *extensa*, 252, 277
— *Gomphrena*, 75
— *Leveillei*, 251
— *Martini*, 251, 277
— *papillosa*, 251
— *papillosa*, 277
— *saligna*, 251, 277
— *Seguini*, 251, 277
— *spelaeicola*, 253, 277
— *subarborea*, 252, 277
— *Vanioti*, 250, 276
— *volkameriaefolia*, 251, 276
- VESTAL, MARY R. and I. W. BAILEY, The orientation of cellulose in the secondary wall of tracheary cells, 185, pl. 206-208, figs.
- and I. W. BAILEY, The significance of certain wood-destroying fungi in the study of the enzymatic hydrolysis of cellulose, 196, pl. 209, 210, figs.
- Viburnum ajugifolium*, 250
— *Cavaleriei*, 250
— *foetidum*, 250
— *foetidum ceanothoides*, 250
— *Valbrayi*, 250
- Vitis cantoniensis*, 71
— *Chaffanjoni*, 220
— *Gentiliana*, 219
— *heterophylla aconitifolia*, 220
- Vitis oligocarpa*, 220
— *Potentilla glabra*, 219
— *Seguini*, 276
- Wallich, Nathaniel. Tentamen florae napalensis illustratae. — Bibliographical notes, 260
- Webera Cavaleriei*, 248
— *Henryi*, 248
— *pallida*, 248
- Wendlandia Dunniana*, 247
— *uvariifolia Dunniana*, 247
- Wikstroemia Bodinieri*, 235, 361
— *Hemsleyana*, 235
- Wisteria dubia*, 68
- Wood-destroying fungi in the study of the enzymatic hydrolysis of cellulose, The significance of certain, 196, pl. 209, 210, figs.
- Wrightia Schlechteri*, 238, 361
— *spec.*, 239
- Xolisma ovalifolia*, 231
— — *lanceolata*, 231
— *villosa pubescens*, 231
- Zanthoxylum Chaffanjoni*, 210
— *myriacanthum*, 209
— *odoratum*, 209
— *rhetoides*, 209
— *trifoliatum*, 71
- Zenobia cerasiflora*, 230
- Zizyphus pubinervis*, 218